

# PROXIMAL HAMSTRING AVULSION REPAIR CLINICAL PRACTICE GUIDELINE

## Background

The posterior thigh consists of three hamstring muscles: semimembranosus, semitendinosus, and biceps femoris. Proximal hamstring injuries and partial tears are common sports injuries; however, complete avulsion tears are more rare. Complete hamstring avulsion injuries occur most often from a slip or a fall that results in forced hip flexion along with knee extension. These injuries result in either complete or partial loss of hamstring function, depending on the severity of the injury and often times require surgical repair due to difficulty with ADL's and sport activities. Surgical repair is required when there is a complete 3-tendon tear or if there is significant retraction (> 2 cm) of two tendons. The surgical repair for a hamstring avulsion is an open procedure that requires anchors placed in the ischium, and sutures from the anchors are then used to secure the tendon to the ischium. If an avulsion fracture of the ischium is involved, it may require fixation via hardware and screws. Following surgery, patients are weight bearing as tolerated and use crutches for the first 1-2 weeks in order to protect the repair.

## Disclaimer

Progression is time and criterion-based, dependent on soft tissue healing, patient demographics, and clinician evaluation. If you are working with an Ohio State Sports Medicine patient and questions arise, please contact the author by calling our office at (614) 293-2385.



**THE OHIO STATE UNIVERSITY**

WEXNER MEDICAL CENTER

For OSUWMC USE ONLY. To license, please contact the OSU Technology Commercialization Office at <https://tco.osu.edu>.

## Summary of Recommendations

<b>General</b>	<ul style="list-style-type: none"> <li>• No end-range hamstring stretching and long-sitting for 6 weeks</li> <li>• No isolated isotonic hamstring strengthening for 8 weeks</li> <li>• Avoid fast walking for 6 weeks</li> </ul>
<b>Weight Bearing</b>	<ul style="list-style-type: none"> <li>• Weight bearing as tolerated with crutches for 0-2 weeks</li> <li>• Gradually increase weight bearing as able and wean off of crutches thereafter</li> </ul>
<b>Patient Reported Outcomes (PRO)</b>	<ul style="list-style-type: none"> <li>• Lower Extremity Functional Scale (LEFS) at each visit</li> <li>• Consider collecting the Hip Outcome Score (HOS) at 1st visit, monthly, and discharge             <ul style="list-style-type: none"> <li>◦ ADL (17 items)   Sports (9 items)</li> </ul> </li> </ul>
<b>Criteria to Discharge Assistive Device</b>	<ul style="list-style-type: none"> <li>• Normalized gait with no increase in pain</li> <li>• Good quad and glute activation and strength</li> <li>• SLR flexion without quad lag</li> </ul>
<b>Criteria to Initiate Plyometric Program</b>	<ul style="list-style-type: none"> <li>• Full, functional, pain free ROM</li> <li>• &gt; 80% quadriceps, hamstring (hip at 0° and 90° of flexion), and hip (using hand-held dynamometer) strength compared to uninjured leg</li> <li>• Squat &gt; 150% body weight (barbell squat or leg press)</li> <li>• 10 forward and lateral step downs from 8" step with proper mechanics</li> </ul>
<b>Criteria to Initiate Running Program</b>	<ul style="list-style-type: none"> <li>• Full, functional, pain-free ROM</li> <li>• &gt; 80% quadriceps, hamstring (hip at 0° and 90° of flexion), and hip (using hand-held dynamometer) strength compared to uninjured leg</li> <li>• Squat &gt; 150% BW (barbell squat or leg press)</li> <li>• 10 forward and lateral step downs from 8" step with proper mechanics</li> <li>• Hop and hold with proper mechanics (uninvolved → involved)</li> <li>• Ability to tolerate 200-250 plyometric foot contacts without reactive pain</li> <li>• No gross visual asymmetry with treadmill/over ground running</li> </ul>
<b>Criteria for Return to Sport</b>	<ul style="list-style-type: none"> <li>• Functional testing:             <ul style="list-style-type: none"> <li>◦ &gt;90% Limb Symmetry Index (LSI) with hop testing</li> </ul> </li> <li>• Isokinetic testing (or hand held dynamometer)             <ul style="list-style-type: none"> <li>◦ &gt;90% strength LSI at 60°/sec, 180°/sec, and 300°/sec testing</li> <li>◦ Hamstring to quadriceps strength ratio of 55-65% bilaterally</li> </ul> </li> <li>• Strength: &gt; 90% body weight with SL leg press</li> <li>• Functional Performance: to date, no return-to-sport criteria have been tested and published for patients undergoing proximal hamstring avulsion repair             <ul style="list-style-type: none"> <li>◦ Patients participating in sports activities should complete a number of sport specific tasks prior to being allowed to return to sport.</li> </ul> </li> <li>• PROs: Score ≥ 90%</li> <li>• No increase in symptoms with sport-specific progression or testing</li> <li>• Physician clearance</li> </ul>



## Phase I: Early Post-Operative Protective Phase (0-4 weeks)

<b>Goals</b>	<ul style="list-style-type: none"> <li>● Protect repair</li> <li>● Pain control</li> <li>● Wean off crutches</li> </ul>
<b>Precautions</b>	<ul style="list-style-type: none"> <li>● Weight bearing status             <ul style="list-style-type: none"> <li>○ Weight-bearing as tolerated (WBAT) with crutches for 0-2 weeks</li> <li>○ Gradually increase weight bearing as able and wean off of crutches thereafter</li> <li>○ Educate patient on taking smaller steps when weaning off of crutches</li> </ul> </li> <li>● Avoid combined hip flexion and knee extension (lengthened hamstring position)</li> <li>● No fast walking for 6 weeks</li> </ul>
<b>Therapeutic Exercises</b>	<ul style="list-style-type: none"> <li>● Glute sets</li> <li>● Ankle pumps</li> <li>● Quad sets</li> <li>● Abdominal bracing</li> <li>● Straight leg raise abduction only</li> <li>● Compression and cryotherapy</li> <li>● Gait training with crutches             <ul style="list-style-type: none"> <li>○ Shortened step length for involved lower extremity</li> </ul> </li> <li>● Gentle PROM of hip, knee and ankle             <ul style="list-style-type: none"> <li>○ Avoid combined hip flexion and knee extension (lengthened hamstring position)</li> </ul> </li> <li>● Gentle soft-tissue mobilization at proximal insertion/incision site once wound is fully closed</li> <li>● Initiate closed-chain terminal knee extensions             <ul style="list-style-type: none"> <li>○ TKE's (resisted quad sets)</li> </ul> </li> </ul>
<b>Criteria to Progress to Phase II</b>	<ul style="list-style-type: none"> <li>● Full hip, knee and ankle PROM in protected positions             <ul style="list-style-type: none"> <li>○ Avoid lengthened hamstring position</li> </ul> </li> <li>● Good quad control in NWB position</li> <li>● Pain and inflammation control</li> <li>● Normalization of gait</li> </ul>



## Phase II: D/C Crutches to Pain free with ADLs (4-6 weeks)

<b>Goals</b>	<ul style="list-style-type: none"> <li>● Protect repair</li> <li>● Initiation of strengthening of hamstring and of proximal and distal musculature (glutes, core, quads)</li> </ul>
<b>Precautions</b>	<ul style="list-style-type: none"> <li>● Avoid end range/aggressive lengthened hamstring position</li> <li>● All exercises to patient tolerance</li> </ul>
<b>Therapeutic Exercises</b>	<ul style="list-style-type: none"> <li>● Initiate stationary bike (no resistance at first)</li> <li>● Weight bearing progression, per patient tolerance</li> <li>● Initiate gentle PROM straight leg hamstring stretching per patient tolerance</li> <li>● Continue soft tissue mobilization</li> <li>● Aquatic activities (if available): forward and retro ambulation, gentle AROM (avoid terminal stretching), gentle partial weight bearing squats</li> <li>● Initiate single leg stance and static proprioceptive activities</li> <li>● Initiate sub-maximal hamstring isometrics in supine             <ul style="list-style-type: none"> <li>○ Avoid lengthened hamstring positions initially</li> <li>○ Begin at 30°, 45°, 60°, 90° knee flexion</li> </ul> </li> <li>● Straight leg raises in flexion (0° to 30° maximum ROM)</li> <li>● Hip strengthening             <ul style="list-style-type: none"> <li>○ SLR abduction and adduction, clamshells, hooklying hip abd isometrics, bridges</li> </ul> </li> <li>● Calf raises</li> <li>● Core strengthening (lumbopelvic control)             <ul style="list-style-type: none"> <li>○ Pelvic tilts, transverse abdominis activation, prone and side planks</li> </ul> </li> </ul>
<b>Criteria to Progress to Phase III</b>	<ul style="list-style-type: none"> <li>● SLR without quad lag</li> <li>● Achieve 45° SLR PROM</li> </ul>



**THE OHIO STATE UNIVERSITY**

WEXNER MEDICAL CENTER

For OSUWMC USE ONLY. To license, please contact the OSU Technology Commercialization Office at <https://tco.osu.edu>.

## Phase III: Pain free ADLs to Initiating Impact Activities (6-12 weeks)

<b>Goals</b>	<ul style="list-style-type: none"> <li>• Correct compensatory movement patterns with functional movements</li> <li>• Optimal neuromuscular control, balance, and proprioception</li> <li>• Normalize strength of core/trunk and glutes</li> <li>• Tolerate single limb support, progressing from single to multi-planar movements</li> <li>• Increase volume/intensity of aerobic activities; restore non-impact cardiovascular fitness</li> <li>• Initiate progressive plyometric activities</li> <li>• Straight leg raise 0°-70° PROM initially and progress to full PROM</li> </ul>
<b>Precautions</b>	<ul style="list-style-type: none"> <li>• Avoid aggressive stretching, especially at end-range</li> <li>• All exercises per patient tolerance</li> </ul>
<b>Criteria to Initiate Plyometrics (Late phase)</b>	<ul style="list-style-type: none"> <li>• Full, functional, pain free ROM</li> <li>• &gt; 80% quadriceps, hamstring (hip at 0° and 90° of flexion), and hip (using hand-held dynamometer) strength compared to uninvolved leg</li> <li>• Squat &gt; 150% body weight (barbell squat or leg press)</li> <li>• 10 forward and lateral step downs from 8" step with proper mechanics</li> </ul>
<b>Therapeutic Exercises</b>	<p><u>Early Exercises:</u></p> <ul style="list-style-type: none"> <li>• Initiate terminal/end-range hamstring stretching</li> <li>• Initiate elliptical, treadmill walking</li> <li>• Initiate gentle isotonic resistive hamstring exercises             <ul style="list-style-type: none"> <li>○ Supine heel slides</li> <li>○ Bilateral hamstring curl machine (light resistance)</li> <li>○ Begin with mid-range strengthening initially</li> </ul> </li> <li>• Bridge progressions</li> <li>• Progress dynamic core strength and trunk stabilization</li> <li>• Progress proprioceptive exercises             <ul style="list-style-type: none"> <li>○ Single leg stance variations with perturbations</li> </ul> </li> <li>• Progressive lower extremity strengthening</li> </ul> <p><u>Late Exercises:</u></p> <ul style="list-style-type: none"> <li>• Initiate partial weight bearing plyometrics on shuttle             <ul style="list-style-type: none"> <li>○ progress to FWBing hop-downs (light)                 <ul style="list-style-type: none"> <li>▪ Start with 2 inch height box/step and progressively increase height</li> <li>▪ Progress from DL to SL</li> </ul> </li> </ul> </li> <li>• Progress full hamstring and quad strengthening program             <ul style="list-style-type: none"> <li>○ Progress bilateral hamstring to unilateral</li> <li>○ Leg press</li> <li>○ Bridging on swiss ball</li> <li>○ Lunges (forward and retro)</li> <li>○ RDLs (DL →SL)</li> <li>○ KB swings</li> </ul> </li> <li>• Advanced core and trunk stabilization exercises</li> <li>• Dynamic proprioceptive activities</li> </ul>



	<ul style="list-style-type: none"> <li>○ Single leg balance activities on BOSU</li> <li>● Resisted ambulation in all directions (ie: cable column, resistance bands, sport cord) <ul style="list-style-type: none"> <li>○ Caution with resisted forward ambulation due to increased hamstring activation</li> </ul> </li> </ul>
<b>Criteria to Progress to Phase IV</b>	<ul style="list-style-type: none"> <li>● Full range of motion and 5/5 MMT <ul style="list-style-type: none"> <li>○ &gt; 80% quadriceps, hamstring (hip at 0° and 90° of flexion), and hip (using hand-held dynamometer) strength compared to uninvolved leg</li> <li>○ Can perform isokinetic testing at 12 weeks to determine readiness to jog</li> </ul> </li> <li>● SLR range of motion within normal limits</li> <li>● Normalized closed chain proprioceptive stability without symptom increase</li> <li>● Tolerate hop downs with appropriate mechanics and no evidence of dynamic instability</li> <li>● Squat &gt; 150% body weight (barbell squat or leg press)</li> <li>● 10 forward and lateral step downs from 8" step with proper mechanics</li> </ul>

## Phase IV: Return to Full Impact/Running (12-16 weeks)

<b>Goals</b>	<ul style="list-style-type: none"> <li>● Progression of full-weight bearing plyometric activities without an increase in symptoms</li> <li>● Isokinetic and functional hop testing for those needing return to sport clearance</li> </ul>
<b>Therapeutic Exercises</b>	<ul style="list-style-type: none"> <li>● Multi-directional plyometrics: DL → SL</li> <li>● Progress single leg strengthening</li> <li>● Progress to eccentric hamstring strengthening</li> </ul>
<b>Criteria to Initiate Running</b>	<ul style="list-style-type: none"> <li>● Full, functional, pain-free ROM</li> <li>● &gt; 80% quadriceps, hamstring (hip at 0° and 90° of flexion), and hip (using hand-held dynamometer or isokinetic testing for quadriceps/hamstring) strength compared to uninvolved leg</li> <li>● Squat &gt; 150% BW (barbell squat or leg press)</li> <li>● 10 forward and lateral step downs from 8" step with proper mechanics</li> <li>● Hop and hold with proper mechanics (uninvolved → involved)</li> <li>● Ability to tolerate 200-250 plyometric foot contacts without reactive pain</li> <li>● No gross visual asymmetry and rhythmic strike pattern with treadmill or over ground running</li> </ul>
<b>Criteria to Progress to V</b>	<ul style="list-style-type: none"> <li>● Jog on treadmill and even surfaces with symmetrical mechanics and no symptoms</li> <li>● Isokinetic testing: &gt;80% LSI</li> <li>● Functional hop testing: &gt; 80% LSI (Appendix A)</li> <li>● Y balance 85% of greater on anterior and posterior lateral reach (Appendix B)</li> </ul>



**THE OHIO STATE UNIVERSITY**

WEXNER MEDICAL CENTER

For OSUWMC USE ONLY. To license, please contact the OSU Technology Commercialization Office at <https://tco.osu.edu>.

## Phase V: Return to Sport/Full Activity (16-24 weeks)

<b>Goals</b>	<ul style="list-style-type: none"><li>● Progressing multi-directional and advanced plyometrics</li><li>● Sport specific drills</li><li>● Physician clearance for return to sport</li></ul>
<b>Therapeutic Exercises</b>	<ul style="list-style-type: none"><li>● Hops to/from BOSU, multidirectional</li><li>● Multidirectional running</li><li>● Resisted forwards running</li><li>● Initiating sport specific drills<ul style="list-style-type: none"><li>○ progress 50% to 75% to full speed/effort</li></ul></li></ul>
<b>Criteria for Discharge and Return to Sports</b>	<ul style="list-style-type: none"><li>● Physician clearance</li><li>● Isokinetic testing: &gt;90% LSI<ul style="list-style-type: none"><li>○ Hamstring to quadricep strength ratio of 55-65% bilaterally</li></ul></li><li>● Functional hop testing: &gt;90% LSI (see appendix)</li><li>● No increase in symptoms with sport specific progressions or testing</li></ul>




**THE OHIO STATE UNIVERSITY**

WEXNER MEDICAL CENTER

For OSUWMC USE ONLY. To license, please contact the OSU Technology Commercialization Office at <https://tco.osu.edu>.

## Appendix A: Single Leg Hop Series

- 1) **Single hop for distance:** Have the subject line their heel up with the zero mark of the tape measure, wearing athletic shoes. The subject then hops as far as he/she can, landing on the same push off leg, for at least 3 seconds. The arms are allowed to move freely during the testing. Allow him/her to perform 2 practice hops on each leg. Then, have the subject perform 2 testing trial, recording each distance from the starting point to the back of the heel. Average the distanced hopped for each limb. The Limb Symmetry Index: Involved limb distance/Uninvolved limb distance X 100%.
- 2) **Cross-over hop for distance:** The subject lines their heel up with the zero mark of the tape measure and hops 3 times on one foot, crossing fully over the center line each time. Each subject should hop as far forward as he/she can on each hop, but only the total distance hopped is recorded. The arms are allowed to move freely during the testing. Allow him/her to perform 2 practice hops on each leg. Then, have the subject perform 2 testing trial, recording each distance from the starting point to the back of the heel. Average the distanced hopped for each limb. The Limb Symmetry Index: Involved limb distance/Uninvolved limb distance X 100%.
- 3) **Triple hop for distance:** The subject lines their heel up with the zero mark of the tape measure and hops 3 times on one foot. Each subject should hop as far forward as he/she can on each hop, but only the total distance hopped is recorded. The arms are allowed to move freely during the testing. Allow him/her to perform 2 practice hops on each leg. Then, have the subject perform 2 testing trial, recording each distance from the starting point to the back of the heel. Average the distanced hopped for each limb. The Limb Symmetry Index: Involved limb distance/Uninvolved limb distance X 100%.
- 4) **Timed 6-meter hop:** The subject lines their heel up at the zero mark of the tape measure and hops, on cue with the tester, as fast as they can the length of the 6-meter tape. The arms are allowed to move freely during the testing. Allow him/her to perform 2 practice hops on each leg. Then, have the subject perform 2 testing trial, recording each distance from the starting point to the back of the heel. Average the distanced hopped for each limb. The Limb Symmetry Index: Involved limb time/Uninvolved limb time X 100%.




**THE OHIO STATE UNIVERSITY**

WEXNER MEDICAL CENTER

For OSUWMC USE ONLY. To license, please contact the OSU Technology Commercialization Office at <https://tco.osu.edu>.



## Appendix B: Y Balance



	Left	Right	Difference
Anterior			
Posteromedial			
Posterolateral			

*Difference should be less than 4 cm for return to sport and preparticipation screening.*

$$\text{Composite Score} = \frac{(\text{Anterior} + \text{Posteromedial} + \text{Posterolateral})}{(3 \times \text{Limb Length})} \times 100$$

Right	
Left	



**THE OHIO STATE UNIVERSITY**

WEXNER MEDICAL CENTER

For OSUWMC USE ONLY. To license, please contact the OSU Technology Commercialization Office at <https://tco.osu.edu>.

**Authors:** Sarah Depp PT, DPT, OCS; Chelseana Lahman, PT, DPT, SCS

**Reviewers:** Jesse Anderson, PT, DPT; Chase Pendley, PT, DPT

**Completion date:** 12/6/2019

## References

- 1) Arner J, Freiman H, Mauro C, Bradley J: Functional Results and Outcomes After Repair of Partial Proximal Hamstring Avulsions at Midterm Follow-up. *Am J of Sports Med* (2019);20:10 1-8
- 2) Belk J et al: Return to Sport After Proximal Hamstring Tendon Repair. *Ortho J of Sports Med* (2019); 6:1-6
- 3) Bourne M et al: An Evidence-Based Framework for Strengthening Exercises to Prevent Hamstring Injury. *Sports Med* (2018);48:251-267
- 4) Brueker PU, Imhoff AB: Functional assessment after acute and chronic complete ruptures of the proximal hamstring tendons. *Knee Surg Sports Traumatol Arthrosc* (2005);13: 411-418
- 5) Clanton TO, Coupe KJ: Hamstring Strains in Athletes: Diagnosis and Treatment. *J Am Acad Orthop Surg* (1998); 6:237-248
- 6) Colosimo AJ, Wyatt HM, Frank KA, Mangine RE: Hamstring Avulsion Injuries. *Oper Tech Sports Med* (2005); 13:80-88
- 7) Kirkland A, Garrison C, Singleton S, Rodrigo J, Boettner F, Stuckey S: Surgical and Therapeutic Management of a Complete Proximal Hamstring Avulsion After Failed Conservative Approach. *J of Orth Sp Phys Ther* (2008); 38;12:754-760
- 8) Leger-St-Jean B, Gorica Z, Magnussen RA, Vasileff WK, Kaeding CC. Accelerated rehabilitation results in good outcomes following acute repair of proximal hamstring ruptures. *Knee Surg Sports Traumatol Arthrosc*. 2019;27:3121-3124
- 9) Lempainen L, Sarimo J, Heikkila J, Mattila K, Orava S: Surgical treatment of partial tears of the proximal origin of the hamstring muscles. *Br J Sports Med* (2006); 688-691
- 10) Moatshe G et al: Repair of Proximal Hamstring Tears: A Surgical Technique. *Arthroscopic Techniques* (2017); 6:2:311-317
- 11) O Mohamed et al: Relationship between wire EMG activity, muscle length, and torque of the hamstrings. *Clinical Biomechanics* (2002); 17:569-579
- 12) Sherry MA, Best TM: A comparison of 2 rehabilitation programs in the treatment of acute hamstring strains. *J of Orth Sp Phys Ther* (2004); 34:3:116-125



**THE OHIO STATE UNIVERSITY**

WEXNER MEDICAL CENTER

For OSUWMC USE ONLY. To license, please contact the OSU Technology Commercialization Office at <https://tco.osu.edu>.