



**Longitudinal Group Review of Systems**

**Gastrointestinal**

- Abdominal pain
- Black tarry stool (melena)
- Bloating
- Bright red blood in the stool (hematochezia)
- Change in stool caliber or color
- Constipation
- Diarrhea
- Difficulty swallowing (dysphagia)
- Early satiety
- Heartburn
- Involuntary loss of stool (fecal incontinence)
- Loss of appetite (anorexia)
- Nausea
- Painful swallowing (odynophagia)
- Rectal pain
- Regurgitation
- Urge to defecate when bowels are empty (tenesmus)
- Vomiting
- Vomiting blood (hematemesis)
- Yellowing of the skin or eyes (jaundice)

**Urinary**

- Absent urine production (anuria)
- Blood in the urine (hematuria)
- Change in urine appearance or odor
- Delay starting urination (urinary hesitancy)
- Decreased urine production (oliguria)
- Dribbling urine
- Flank pain
- Foamy urine
- Frequent urination (urinary frequency)
- Frequent urination at night (nocturia)
- Incomplete bladder emptying (urinary retention)
- Increased urine production (polyuria)
- Intermittent flow of urinary stream (urinary intermittency)
- Involuntary loss of urine (urinary incontinence)
- Painful urination (dysuria)
- Straining to urinate
- Suprapubic pain
- Swelling of the hands, feet, or face (edema)
- Urge to urinate (urinary urgency)
- Weak urinary stream